New Mexico

Population

- 0.7% of US population
- 1.7% blacks/African Americans, 46.3% Hispanics/Latinos, 40.5% whites, and 11.5% other races/ethnicities

HIV Diagnoses

- 149 diagnoses, 2013
 - $\circ~~0.3\%$ of diagnoses made in the US, 2013
- 2,558 persons living with diagnosed HIV infection, 2012
 - $^{\circ}$ $\,$ 0.3% of persons living with diagnosed HIV infection in the US, 2012

Progress toward HIV Prevention and Care Objectives

| Indicator: Objective | National 2015 Goal | Year 1 Baseline | Year 2 | Year 3 | Improved Year 3 vs. Year 1 | State Ranking Year 3 | Met 2015 Goal in Year 3 |
|---|--------------------------|--------------------|----------|---------|----------------------------------|----------------------------|-------------------------------|
| PREVENT NEW HIV INFECTIONS | | | | | | | |
| Increase HIV testing (ever tested) by 4%, 2011-2013 | 44.2 | 39.1 | 38.8 | 41.1 | | 27 of 51 | |
| Increase knowledge of HIV status to 90%, 2010-2012 | 90.0 | 86 | 88 | 89 | | 6 of 42 | |
| Reduce late-stage HIV diagnosis by 25%, 2011-2013 | 19.1 | 23.9 | 28.1 | 22.2 | \checkmark | 10 of 51 | |
| INCREASE ACCESS TO HI | V CARE ANI | D IMPRO | VE HEAL | тн оитс | OMES | | |
| Increase linkage to HIV medical care to 85%, 2011-2013 | 85.0 | NA | NA | NA | | - of 28 | |
| Increase retention in HIV medical care by 10%, 2010-2012 | 56.0 | NA | NA | NA | | - of 28 | |
| Increase viral suppression among persons in HIV medical care by 10%, 2010-2012 | 80.0 | NA | NA | NA | | - of 28 | |
| Decrease the death rate among HIV-diagnosed persons (per 1,000 persons) by 10%, 2010-2012 | 21.7 | 24.9 | 22.8 | 20.8 | ✓ | 35 of 51 | ✓ |
| REDUCE HIV-RELATED | DISPARITI | ES AND I | HEALTH I | INEQUIT | IES | | |
| Increase HIV testing (ever tested) among race/ethnicity gro | oups by 4%, 2 | 2011-2013 | | | | | |
| Blacks/African Americans | 68.8 | 53.0 | 70.0 | 58.2 | | 33 of 40 | |
| Hispanics/Latinos | 46.6 | 36.2 | 34.0 | 39.4 | | 34 of 39 | |
| Whites | 39.6 | 42.9 | 42.6 | 43.2 | | 10 of 51 | \checkmark |
| Persons of other races/ethnicities | 40.7 | 34.6 | 41.5 | 38.1 | | 31 of 39 | |
| Reduce late-stage HIV diagnosis among race/ethnicity group | ups by 25%, | 2011-2013 | 3 | | | | |
| Blacks/African Americans | 17.6 | 14.3 | 0.0 | 0.0 | | 1 of 51 | \checkmark |
| Hispanics/Latinos | 21.6 | 28.6 | 33.8 | 22.5 | \checkmark | 14 of 51 | |
| Whites | 19.5 | 20.6 | 20.7 | 29.4 | | 35 of 51 | |
| Persons of other races/ethnicities | 20.4 | 18.5 | 22.7 | 17.4 | \checkmark | 9 of 49 | \checkmark |

Note. For information on indicator definitions and data sources, see <u>Technical Notes</u>. Abbreviation: PLWH, persons living with diagnosed HIV infection; NA, not available.

 $[\]checkmark$ = Yes: improved since baseline (chi-square p<0.05 or >5% change); met 2015 goal.